

Published in final edited form as:

Gut. 2009 January ; 58(1): 90–96. doi:10.1136/gut.2008.155473.

CpG island methylator phenotype, microsatellite instability, *BRAF* mutation and clinical outcome in colon cancer

Shuji Ogino^{*}, Katsuhiko Noshō, Gregory J. Kirkner, Takako Kawasaki, Jeffrey A. Meyerhardt, Massimo Loda, Edward L. Giovannucci, and Charles S. Fuchs

From Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA (SO, NK, TK, JAM, ML, CSF); Department of Pathology, Brigham and Women's Hospital, Boston, and Harvard Medical School, Boston, MA (SO, ML); Department of Epidemiology, Harvard School of Public Health, Boston, MA (SO, ELG); Channing Laboratory, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA (GJK, ELG, CSF)

Abstract

Background—The CpG island methylator phenotype (CIMP) characterized by widespread promoter methylation is associated with microsatellite instability (MSI) and *BRAF* mutation in colorectal cancer. The independent effect of CIMP, MSI and *BRAF* mutation on patient outcome remains uncertain.

Methods—Utilizing 649 colon cancers (stage I–IV) in two independent cohort studies, we quantified DNA methylation in 8 CIMP-specific promoters [*CACNA1G*, *CDKN2A* (p16), *CRABP1*, *IGF2*, *MLH1*, *NEUROG1*, *RUNX3*, and *SOCS1*], as well as *MINT1*, *MINT31*, p14, *HIC1*, *IGFBP3*, *MGMT* and *WRN* by MethyLight. We examined MSI, *KRAS* and *BRAF* status. Cox proportional hazard models computed hazard ratios (HRs) for colon cancer-specific and overall mortalities, adjusting for patient characteristics and tumoral molecular features.

Results—After adjustment for other predictors of patient survival, patients with CIMP-high cancers [126 (19%) tumors with $\geq 6/8$ methylated CIMP-specific promoters] experienced a significantly low colon cancer-specific mortality [multivariate HR 0.44, 95% confidence interval (CI) 0.22–0.88], whereas *BRAF* mutation was significantly associated with a high cancer-specific mortality (multivariate HR 1.97, 95% CI, 1.13–3.42). A trend toward a low cancer-specific mortality was observed for MSI-high tumors (multivariate HR 0.70, 95% CI, 0.36–1.37). In stratified analyses, CIMP-high tumors were associated with a significant reduction in colon cancer-specific mortality, regardless of both MSI and *BRAF* status. The relation between CIMP-high and lower mortality appeared to be consistent across all stages. *KRAS* mutation was unrelated to patient outcome.

Conclusion—CIMP-high appears to be an independent predictor of a low colon cancer-specific mortality, while *BRAF* mutation is associated with a high colon cancer-specific mortality.

*Corresponding Author: Shuji Ogino, M.D., Ph.D. Center for Molecular Oncologic Pathology, Dana-Farber Cancer Institute, Brigham and Women's Hospital, Harvard Medical School, 44 Binney Street, Room JF-215C, Boston, MA 02115 USA, Phone: +1-617-632-3978, Fax: +1-617-277-9015, E-mail: E-mail: shuji_ogino@dfci.harvard.edu.

Competing Interest: None to declare.

The manuscript has not been published previously and is not being considered concurrently by any other publication. This manuscript acknowledges all sources of support for the work.

Licence for Publication: The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive licence (or non exclusive for government employees) on a worldwide basis to the BMJ Publishing Group Ltd and its Licensees to permit this article (if accepted) to be published in Gut editions and any other BMJ PGL products to exploit all subsidiary rights, as set out in our licence (<http://gut.bmjournals.com/fora/licence.dtl>).

Keywords

colorectal cancer; CIMP; methylation; MSI; prognosis

INTRODUCTION

Epigenetic aberrations are thought to be an important mechanism in human carcinogenesis. [1,2] A number of tumor suppressor genes are silenced by promoter CpG island methylation in colon cancers.[2,3] A subset of colon cancers exhibit widespread promoter methylation, referred to as the CpG island methylator phenotype (CIMP).[2,4–6] which is associated with microsatellite instability (MSI).[7,8] CIMP-high colon cancers have been associated with older age, cigarette smoking, proximal tumor location, female gender, poor differentiation, *BRAF* mutation, wild-type *TP53*, inactive β -catenin/*WNT* and stable chromosomes,[8–19] and many of these associations are independent of MSI status.[9,16–19]

Among patients with colon cancer, MSI has generally been associated with good prognosis in most,[20] though not all studies.[21] On the other hand, the presence of *BRAF* mutations in tumors has been characteristically associated with an inferior patient survival.[22] In contrast, results for CIMP have been conflicting.[22–28] These inconsistent results likely reflect differences in patient cohorts, methylation markers examined, and the variable inclusion of data on other potentially confounding molecular events, such as MSI and *BRAF* in multivariate analysis models.

We therefore examined both genetic and epigenetic alterations among colon cancer patients participating in two large prospective cohort studies, to assess the independent effect of CIMP, MSI and *BRAF* mutation on patient outcome. Furthermore, to assess CpG island methylation, we utilized quantitative DNA methylation assays (MethyLight technology) on a validated expanded panel of 8 markers that appears to well characterize the presence or absence of CIMP-high in colorectal cancers.[8,29]

METHODS

Study population

We utilized the databases of two independent prospective cohort studies; the Nurses' Health Study (N=121,700 women followed since 1976),[30,31] and the Health Professional Follow-up Study (N=51,500 men followed since 1986).[31] On each biennial follow-up questionnaire, participants were asked whether they had a diagnosis of colon cancer during the previous 2 years. When a participant (or next of kin for decedents) reported colon cancer, we sought permission to obtain medical records. Study physicians, while blinded to exposure data, reviewed all records related to colon cancer, and recorded the date of cancer diagnosis, AJCC (American Joint Committee on Cancer) stage and tumor location. For nonresponders, we searched the National Death Index to discover deaths and ascertain any diagnosis of colon cancer that was a primary cause of death or a secondary diagnosis. Approximately 96% of all incident colon cancer cases were identified through these methods. We collected paraffin-embedded tissue blocks from hospitals where patients underwent resections of primary colon cancers.[31] Tissue sections from all cases in this study were reviewed by a pathologist (S.O.). Tumor grade was categorized as high ($\leq 50\%$ glandular area) or low ($> 50\%$ glandular area). We excluded rectal cancers and cases that were preoperatively treated with radiation and/or chemotherapy. Based on availability of tissue samples, we included a total of 649 colon cancer cases (283 from the men's cohort and 366 from the women's cohort) diagnosed up to 2002. Written informed consent was obtained from all study subjects. This study was approved by

the Human Subjects Committees at Brigham and Women's Hospital and Harvard School of Public Health.

Measurement of mortality

Patients were observed until death or June 2006, whichever came first. Ascertainment of deaths included reporting by the family or postal authorities. The names of persistent nonresponders were searched in the National Death Index. The cause of death was assigned by physicians blinded to information on lifestyle exposures and molecular changes in colon cancer. In rare patients who died as a result of colon cancer not previously reported, we obtained medical records with permission from next of kin. More than 98% of deaths in the cohorts were identified by these methods.

Genomic DNA extraction and sequencing of *KRAS* and *BRAF*

Genomic DNA from paraffin-embedded tissue was extracted, and whole genome amplification was performed by PCR using random 15-mer primers.[32] PCR and sequencing targeted for *KRAS* codons 12 and 13, and *BRAF* codon 600 were performed as previously described.[32, 33]

Real-time PCR (MethyLight) for quantitative DNA methylation analysis

Bisulfite treatment on genomic DNA and subsequent real-time PCR (MethyLight)[34] were validated and performed as previously described.[35] We quantified DNA methylation in 8 CIMP-specific promoters (*CACNA1G*, *CDKN2A (p16)*, *CRABP1*, *IGF2*, *MLH1*, *NEUROG1*, *RUNX3* and *SOCS1*)[8,29] (which were selected from screening of 195 CpG islands in the human genome[8,17]), as well as *HIC1*, *MINT1*, *MINT31*,[36] *MGMT*,[35] *IGFBP3* and *WRN*. [25] The PCR condition was initial denaturation at 95°C for 10 min followed by 45 cycles of 95°C for 15 sec and 60°C for 1 min.

CIMP-high was defined as $\geq 6/8$ methylated markers using the 8-marker CIMP panel, CIMP-low/0 as $\leq 5/8$ methylated markers, and CIMP-0 as 0/8 methylated markers, according to the previously established criteria.[29]

Microsatellite Instability (MSI) Analysis

MSI status was determined using D2S123, D5S346, D17S250, BAT25, BAT26,[37] BAT40, D18S55, D18S56, D18S67 and D18S487 (i.e., 10-marker panel).[29] A "high degree of MSI" (MSI-high) was defined as the presence of instability in $\geq 30\%$ of the markers, and "microsatellite stability" (MSS) as no unstable marker or instability in $< 30\%$ of the markers. When tumors with instability in $< 30\%$ of the markers (i.e., "MSI-low") was compared to tumors with no unstable marker, "MSI-low" did show no prognostic value (data not shown). Thus, we combined "MSI-low" tumors into MSS tumors in further analyses.

Statistical analysis

Cox proportional hazard models were used to calculate hazard ratios (HRs) of death according to molecular features in tumor (i.e., MSI, CIMP and *BRAF* mutation), adjusted for age, sex, year of diagnosis, tumor stage, tumor location, tumor grade, and the molecular variables. In the analyses for colon cancer-specific mortality, death as a result of colon cancer was the primary end point and deaths as a result of other causes were censored. Age and year of diagnosis were used as continuous variables, and all of the other variables were used as categorical variables. When information on tumor location (1.4% missing), tumor stage (7.4% missing), *KRAS* (0.3% missing) or *BRAF* (2.8% missing) was missing, we assigned a separate ("missing") indicator variable and included those cases in the multivariate analysis model. We confirmed that excluding cases with a missing variable did not significantly alter results (data

not shown). An interaction was assessed by including the cross product of two variables of interest in the analysis model. The Kaplan-Meier method was used to describe the distribution of colon cancer-specific and overall survival time, and the log-rank test was performed to test the null hypothesis of no difference in survival time distributions among all subtypes. The chi square test was used to examine an association between categorical variables. The t-test assuming unequal variances was used to compare mean ages. All analyses used SAS version 9.1 (SAS Institute, Cary, NC) and all p values were two-sided.

RESULTS

CpG island methylator phenotype (CIMP) and microsatellite instability (MSI)

Among 649 colon cancer patients with available tissue specimens, there were 281 deaths, including 163 colon cancer-specific deaths. Among all patients, 121 (19%) demonstrated MSI-high; 126 (19%) were CIMP-high ($\geq 6/8$ methylated CIMP-specific markers; [29] i.e., *CACNA1G*, *CDKN2A*, *CRABP1*, *IGF2*, *MLH1*, *NEUROG1*, *RUNX3* and *SOC31*); 238 (37%) demonstrated a mutation in *KRAS*; and 105 (17%) demonstrated a mutation in *BRAF*. Tumors were distributed bimodally according to the number of methylated CIMP markers (Figure 1), and *BRAF* mutations were common in CIMP-high tumors while *KRAS* mutations were common in CIMP-low tumors (1/8–5/8 methylated markers). We assessed baseline patient characteristics according to MSI and CIMP status (Table 1).

MSI-high tumors were more likely to originate in the proximal colon and possess *BRAF* mutations and CIMP-high status, and less likely to present with stage III or IV disease. CIMP-high tumors were also more likely to originate in the proximal colon and possess *BRAF* mutations.

When we jointly classified tumors by MSI and CIMP status, MSS (microsatellite stable) CIMP-high tumors had a greater prevalence of stage IV disease (36%; 13/36; $p=0.0004$, compared to all other subtypes) when compared to MSI-high CIMP-high tumors (4%; 3/84), MSI-high CIMP-low/0 tumors (6%; 2/32) or MSS CIMP-low/0 tumors (14%; 65/452). In addition, *BRAF* mutations were found in 62% (53/86) of MSI-high CIMP-high tumors, 57% (21/37) of MSS CIMP-high tumors, none (0/32) of MSI-high CIMP-low/0 tumors, and 6% (31/476) of MSS CIMP-low/0 tumors.

Molecular features in colon cancer and patient survival

We assessed the influence of MSI, CIMP, and *BRAF* mutation on patient survival, independent of the clinical and other tumoral variables (Table 2).

Compared to patients with MSS tumors, those with MSI-high tumors experienced a significant reduction in colon cancer specific mortality in a univariate analysis [hazard ratio (HR) 0.38, 95% confidence interval (CI), 0.22–0.66]; however, in the multivariate model that adjusted for CIMP, *KRAS*, *BRAF* and patient characteristics, the effect of MSI-high was attenuated. This attenuation in the effect of MSI-high was principally the result of adjusting for tumor stage; when we simply adjusted for tumor stage, the HR for colon cancer-specific mortality for MSI-high tumors was 0.73 (95% CI, 0.42–1.28).

In addition, compared to CIMP-0, CIMP-high tumors were associated with a non-significant reduction in colon cancer specific mortality in a univariate analysis (HR 0.88, 95% CI, 0.57–1.38), which became statistically significant after adjusting for other molecular and patient characteristics (multivariate HR 0.44, 95% CI, 0.22–0.88). The greater beneficial effect of CIMP-high status in the multivariate model was principally the result of adjusting for *BRAF* mutational status; when we simply adjusted for *BRAF* status, the HR for colon cancer-specific mortality for CIMP-high tumors was 0.45 (95% CI, 0.26–0.79).

In both univariate and multivariate analyses, *BRAF* mutation was associated with a significant increase in colon cancer-specific mortality (multivariate HR 1.97, 95% CI, 1.13–3.42). In contrast, *KRAS* mutation was not associated with patient outcome. Of note, the aforementioned molecular events did not significantly influence all-cause mortality.

Although statistical power was diminished for individual patient subsets, the influence of CIMP status on colon cancer-specific mortality appeared similar among patients with either early (I and II) or advanced (III and IV) pathologic stages of disease (p for interaction =0.93). Compared to CIMP-low/0 tumors, the multivariate HR for colon cancer-specific mortality in CIMP-high tumors was consistently low across all stages (I to IV) (Table 3).

In contrast, any apparent effect of CIMP-high on overall mortality was limited to patients with stage III/IV disease; the multivariate HR for all-cause mortality was 1.49 (95% CI, 0.66–3.34) for stage I/II patients and 0.58 (95% CI, 0.29–1.15) for stage III/IV patients.

Next, we examined whether the effect of CIMP or *BRAF* mutation on survival differed between the cohort studies. The effect of CIMP-high did not significantly differ between the male cohort (Health Professionals Follow-up Study) and the female cohort (Nurses' Health Study; p for interaction =0.59). Likewise, the effect of *BRAF* mutation did not significantly differ between the male cohort and the female cohort (p for interaction =0.35).

To eliminate potential confounding effect of HNPCC (hereditary nonpolyposis colorectal cancer), we identified 19 possible or suspected HNPCC cases [i.e., MSI-high CIMP-low/0 tumors (none of which turned out to be *BRAF*-mutated) with any of the followings: (1) positive family history of colorectal cancer in at least one first-degree relative; (2) loss of *MLH1* without evidence of *MLH1* methylation; (3) loss of *PMS2* without evidence of *MLH1* loss; (4) loss of *MSH2* and/or *MSH6*]. After we excluded these 19 cases, multivariate Cox regression analysis showed following results for colon cancer-specific mortality: HR for MSI-high, 0.68 (95% CI, 0.34–1.35); HR for CIMP-high, 0.41 (95% CI, 0.20–0.83); HR for *BRAF* mutation, 1.85 (95% CI, 1.12–3.06). These results were similar to Table 2.

We compared different CIMP panels consisting of different sets of markers. Multivariate HRs for colon cancer-specific mortality in CIMP+ vs. CIMP- were as follows: HR 0.57 (95% CI, 0.32–1.01) by a panel consisting of *CACNA1G*, *IGF2*, *NEUROG1*, *RUNX3* and *SOCS1*;^[8] HR 0.81 (95% CI, 0.55–1.20) by a panel consisting of *CDKN2A* (p16), *MINT1*, *MINT31*, *MLH1* and p14;^[26] HR 0.74 (95% CI, 0.46–1.20) by a panel consisting of *CDKN2A*, *HIC1*, *MINT1*, *MINT31* and *MLH1*; HR 0.54 (95% CI, 0.30–0.99) by a panel consisting of *CACNA1G*, *CDKN2A*, *CRABP1*, *MLH1* and *NEUROG1*;^[17] HR 0.65 (95% CI, 0.37–1.12) by a panel consisting of *CACNA1G*, *CDKN2A*, *CRABP1*, *IGF2*, *MLH1*, *NEUROG1*, *RUNX3*, *SOCS1*, *IGFBP3*, *MINT1*, *MINT31*, *MGMT* and *WRN*.^[25] These results indicate that a variation in methylation markers in CIMP panels can result in a variation in associations with patient outcome, which may, at least in part, explain the discrepancy of different studies on CIMP and patient outcome. In the current study, we utilized the validated 8-marker panel in light of our prior published work.^[29]

Combined MSI/CIMP status and patient survival

We further stratified patients according to both MSI and CIMP status to assess the joint effect on patient outcome (Table 4), because molecular classification based on MSI and CIMP status is increasingly important.^[38,39]

Compared to patients whose tumors were both MSS and CIMP-low/0, those with CIMP-high tumors experienced a significant reduction in colon cancer-specific mortality, regardless of MSI status. A combination of MSI and CIMP determinations might differentiate ~24% [(38

+33+88)/649] of tumors (either CIMP-high or MSI-high) with good prognosis (HR estimates 0.17–0.40) from the other ~76% of tumors (MSS CIMP-low/0).

Combined MSI/*BRAF* status and patient survival

Similarly, we stratified patients according to both MSI and *BRAF* status to assess joint effect on patient outcome. Compared to patients whose tumors were both MSS and *BRAF*-mutated, those with MSI-high/*BRAF*-wild-type tumors showed a significant reduction in colon cancer-specific mortality (Table 4). Notably, there was no protective effect of MSI-high among *BRAF*-mutated tumors; compared to MSS *BRAF*-mutated tumors, the multivariate HR for colon cancer-specific mortality among MSI-high *BRAF*-mutated tumors was 1.09 (95% CI, 0.48–2.51).

Combined CIMP/*BRAF* status and patient survival

We also stratified patients according to both CIMP and *BRAF* status to assess the joint effect on patient outcome. Colon cancer-specific survival at 5 years was 45% for patients with CIMP-low/0 *BRAF*-mutated tumors, 80% for CIMP-low/0 *BRAF*-wild-type tumors, 74% for CIMP-high *BRAF*-mutated tumors, and 86% for CIMP-high *BRAF*-wild-type tumors (multi-group log rank $p < 0.0001$; Figure 2A). Similarly, overall survival at 5 years was lower in CIMP-low/0 *BRAF*-mutated tumors than the other subtypes (multi-group log rank $p = 0.0015$; Figure 2B).

In a multivariate analysis, when compared to patients with CIMP-low/0 *BRAF*-mutated tumors, those with CIMP-high tumors demonstrated a significantly lower colon cancer-specific mortality regardless of *BRAF* status (Table 4). Moreover, the adverse effect of *BRAF* mutation on patient survival was not apparent when tumors also demonstrated CIMP-high.

DISCUSSION

In this cohort of patients with colon cancer, we examined the effect of the CpG island methylator phenotype (CIMP), microsatellite instability (MSI), and *BRAF* mutation on patient outcome. CIMP-high status was independently associated with a low cancer-specific mortality, whereas *BRAF* mutation was associated with a significant increase in cancer-specific mortality. Consistent with other studies,[20] we found that MSI-high tumors showed a trend towards an association with longer survival. Of note, the adverse effect of *BRAF* mutation appeared to be limited to tumors that were not CIMP-high. Although our observations need to be confirmed by other independent studies, the associations of CIMP and *BRAF* mutation with clinical outcome were consistent across the two independent prospective cohort studies in this analysis.

The relationship between CIMP, MSI, and *BRAF* mutations in colon cancer is complex. In our cohort, 70% of *BRAF*-mutated tumors exhibited CIMP-high, and 70% of CIMP-high tumors exhibited MSI-high. Among patients who do not manifest hereditary nonpolyposis colorectal cancer (HNPCC), MSI-high is often the consequence of promoter methylation (and subsequent silencing) of *MLH1*, a DNA mismatch repair gene.[8] In fact, CIMP and *BRAF* tests are used to exclude HNPCC among patients who exhibit MSI-high, since HNPCC seldom exhibits CIMP or *BRAF* mutation.[8,40,41]

Studying epigenetic and/or genetic alterations is increasingly important in cancer research.[3, 42–44] To decipher the apparently complex effect of CIMP and *BRAF* mutation on patient survival, we utilized a validated expanded panel of 8 methylation markers for CIMP diagnosis in colorectal cancer.[8,29] To determine DNA methylation status at each locus, we used a quantitative method that appears to reproducibly differentiate high-level from low-level methylation.[35] Our validated criteria for CIMP-high are based on the bimodal distribution of tumors according to the number of methylated CIMP markers, and the observation that

CIMP-high is associated with *BRAF* mutation while CIMP-low is associated with *KRAS* mutation.[33,45] Our large sample size from the two independent cohort studies enabled us to estimate the frequencies of specific molecular features (e.g., CIMP-high, etc.) and cancer death rates at the population level.

Although prognostic factors have been extensively investigated for colon cancer,[20–22,46–48] previous studies of CIMP and survival in colon cancer have yielded somewhat inconsistent results.[22–28] Some studies suggested an adverse effect of CIMP on survival of patients with MSS tumors.[23,25,26] However, accumulating evidence has been suggested that MSI-high tumors are associated with good prognosis regardless of CIMP status,[22,23] which is in agreement with our current study (Table 4). *BRAF* mutation has been associated with worse survival in MSS tumors, but there was little prognostic value of CIMP in multivariate analysis.[22,27] Our findings of good prognosis in CIMP-high tumors appear to differ from the data in the previous studies.[23,25,26] These discrepant observations might have resulted from differences in patient cohorts, methylation markers, criteria for CIMP, and/or the variable inclusion of other potential confounders (such as *BRAF* mutation) in multivariate analysis models. In particular, we have previously observed worse prognosis associated with CIMP-high tumors in stage IV colorectal cancer in small phase I/II clinical trials.[25] The possible reasons for the discrepant results are as follows: 1) A selection bias in the small clinical trials with only 5 CIMP-high tumors might have caused this discrepancy. 2) Data in our previous study[25] with only 5 CIMP-high tumors might simply be the result by chance in the setting of a small patient population. A p value by the log-rank test is calculated by the Mantel-Haenszel chi-square test that can offer a far more accurate p value with a large sample size. Thus, we would emphasize the importance of a large sample size in any clinical study. Because of the use of the expanded CIMP marker panel (including the 5 new markers described by Weisenberger et al.[8]) in the current study, good prognosis might be specifically associated with CIMP-high tumors defined by these new CIMP makers. Our observations of good prognosis in CIMP-high tumors appeared to be consistent across all stages (I to IV), further supporting that CIMP-high tumor is a biologically indolent subtype. In addition, we found that, after jointly examining CIMP and *BRAF* status, CIMP-high predicted a lower colon cancer-specific mortality (regardless of *BRAF* status) compared to CIMP-low/0 *BRAF*-mutated tumors, whereas the deleterious effect of *BRAF* mutation was not as evident in patients with CIMP-high tumors.

In our cohorts, data on cancer treatment are limited. Nonetheless, it is unlikely that chemotherapy use differed according to tumoral CIMP, MSI or *BRAF* status, especially since such data were not typically available to patients or treating physicians. It still remains a possibility that differential response to chemotherapy according to a specific molecular variable (e.g., MSI) might confound our findings. Further studies are necessary to examine whether response to chemotherapy may be differentially influenced by specific molecular features in colon cancer. In addition, beyond cause of mortality, data on cancer recurrences were not available in these cohorts. Nonetheless, given the median survival for metastatic colon cancer was approximately 10 to 12 months during much of the time period of this study,[49] colon cancer-specific mortality should be a reasonable surrogate for cancer-specific outcomes.

Despite the apparent effects of CIMP, MSI, and *BRAF* mutation on colon cancer-specific mortality, the influence of these tumoral events on overall mortality was markedly attenuated, which may have reflected the inclusion of earlier stage (I and II) patients in our analysis. In fact, when we limited our analysis to patients with either stage III or IV cancer, we observed similar effects of CIMP on both cancer-specific and all-cause mortality. Moreover, when we jointly classified patients according to both CIMP and *BRAF* status, we observed similar trends for cancer-specific and overall mortality among the entire patient cohort (Table 4 and **Figure 4**).

In conclusion, this large prospective study of colon cancer patients suggests that CIMP-high is independently associated with a low cancer-specific mortality. While *BRAF* mutation is associated with worse survival, CIMP-high appears to eliminate the adverse effect of *BRAF* mutation. Finally, while our data confirm the extensive body of evidence supporting a better prognosis for patients with MSI-high tumors, the good prognosis associated with MSI-high was abrogated in the presence of a *BRAF* mutation. Our finding that CIMP-high is an independent predictor of cancer survival may have significant clinical implications, although it needs to be confirmed by additional independent studies. Future studies to validate our observations should consider a joint examination of CIMP, MSI and *BRAF* mutation to decipher the role of these molecular features in biological and clinical behavior of colon cancer.

Acknowledgements

This work was supported by The U.S. National Institute of Health grants P01 CA87969, P01 CA55075, P50 CA127003, and K07 CA122826 (to S.O.), the Bennett Family Fund for Targeted Therapies Research, and the Entertainment Industry Foundation (EIF). K.N. was supported by a fellowship grant from the Japanese Society for the Promotion of Science.

We thank the Nurses' Health Study and Health Professionals Follow-up Study cohort participants who have generously agreed to provide us with biological specimens and information through responses to questionnaires. We thank Walter Willett, Sue Hankinson, and many other staff members who implemented and have maintained the cohort studies. We deeply thank Peter Laird, Daniel Weisenberger and Mihaela Campan for their assistance in the development of the MethyLight assays.

Abbreviations

| | |
|--------------|---|
| AJCC | American Joint Committee on Cancer |
| CI | confidence interval |
| CIMP | CpG island methylator phenotype |
| HNPCC | hereditary nonpolyposis colorectal cancer |
| HPFS | Health Professionals Follow-up Study |
| HR | hazard ratio |
| MSI | microsatellite instability |
| MSS | microsatellite stable |
| NHS | Nurses' Health Study |

References

1. Jones PA, Baylin SB. The epigenomics of cancer. *Cell* 2007;128:683–92. [PubMed: 17320506]
2. Issa JP. CpG island methylator phenotype in cancer. *Nat Rev Cancer* 2004;4:988–93. [PubMed: 15573120]

3. Wong JJ, Hawkins NJ, Ward RL. Colorectal cancer: a model for epigenetic tumorigenesis. *Gut* 2007;56:140–8. [PubMed: 16840508]
4. Toyota M, Ahuja N, Ohe-Toyota M, et al. CpG island methylator phenotype in colorectal cancer. *Proc Natl Acad Sci U S A* 1999;96:8681–6. [PubMed: 10411935]
5. Grady WM. CIMP and colon cancer gets more complicated. *Gut* 2007;56:1498–500. [PubMed: 17938428]
6. Teodoridis JM, Hardie C, Brown R. CpG island methylator phenotype (CIMP) in cancer: Causes and implications. *Cancer Lett* 2008;268:177–86. [PubMed: 18471961]
7. Toyota M, Ohe-Toyota M, Ahuja N, et al. Distinct genetic profiles in colorectal tumors with or without the CpG island methylator phenotype. *Proc Natl Acad Sci U S A* 2000;97:710–5. [PubMed: 10639144]
8. Weisenberger DJ, Siegmund KD, Campan M, et al. CpG island methylator phenotype underlies sporadic microsatellite instability and is tightly associated with BRAF mutation in colorectal cancer. *Nat Genet* 2006;38:787–93. [PubMed: 16804544]
9. Samowitz WS, Albertsen H, Sweeney C, et al. Association of smoking, CpG island methylator phenotype, and V600E BRAF mutations in colon cancer. *J Natl Cancer Inst* 2006;98:1731–8. [PubMed: 17148775]
10. van Rijnsoever M, Grieu F, Elsaleh H, et al. Characterisation of colorectal cancers showing hypermethylation at multiple CpG islands. *Gut* 2002;51:797–802. [PubMed: 12427779]
11. Whitehall VL, Wynter CV, Walsh MD, et al. Morphological and molecular heterogeneity within nonmicrosatellite instability-high colorectal cancer. *Cancer Res* 2002;62:6011–4. [PubMed: 12414620]
12. Hawkins N, Norrie M, Cheong K, et al. CpG island methylation in sporadic colorectal cancers and its relationship to microsatellite instability. *Gastroenterology* 2002;122:1376–87. [PubMed: 11984524]
13. Kambara T, Simms LA, Whitehall VLJ, et al. BRAF mutation is associated with DNA methylation in serrated polyps and cancers of the colorectum. *Gut* 2004;53:1137–44. [PubMed: 15247181]
14. Nagasaka T, Sasamoto H, Notohara K, et al. Colorectal cancer with mutation in BRAF, KRAS, and wild-type with respect to both oncogenes showing different patterns of DNA methylation. *J Clin Oncol* 2004;22:4584–94. [PubMed: 15542810]
15. Goel A, Nagasaka T, Arnold CN, et al. The CpG Island Methylator Phenotype and Chromosomal Instability Are Inversely Correlated in Sporadic Colorectal Cancer. *Gastroenterology* 2007;132:127–38. [PubMed: 17087942]
16. Samowitz W, Albertsen H, Herrick J, et al. Evaluation of a large, population-based sample supports a CpG island methylator phenotype in colon cancer. *Gastroenterology* 2005;129:837–45. [PubMed: 16143123]
17. Ogino S, Cantor M, Kawasaki T, et al. CpG island methylator phenotype (CIMP) of colorectal cancer is best characterised by quantitative DNA methylation analysis and prospective cohort studies. *Gut* 2006;55:1000–6. [PubMed: 16407376]
18. Kawasaki T, Nosho K, Ohnishi M, et al. Correlation of beta-catenin localization with cyclooxygenase-2 expression and CpG island methylator phenotype (CIMP) in colorectal cancer. *Neoplasia* 2007;9:569–77. [PubMed: 17710160]
19. Ogino S, Kawasaki T, Kirkner GJ, et al. 18q loss of heterozygosity in microsatellite stable colorectal cancer is correlated with CpG island methylator phenotype-negative (CIMP-0) and inversely with CIMP-low and CIMP-high. *BMC Cancer* 2007;7:72. [PubMed: 17474983]
20. Popat S, Hubner R, Houlston RS. Systematic Review of Microsatellite Instability and Colorectal Cancer Prognosis. *J Clin Oncol* 2005;23:609–18. [PubMed: 15659508]
21. Kim GP, Colangelo LH, Wieand HS, et al. Prognostic and predictive roles of high-degree microsatellite instability in colon cancer: a National Cancer Institute-National Surgical Adjuvant Breast and Bowel Project Collaborative Study. *J Clin Oncol* 2007;25:767–72. [PubMed: 17228023]
22. Samowitz WS, Sweeney C, Herrick J, et al. Poor survival associated with the BRAF V600E mutation in microsatellite-stable colon cancers. *Cancer Res* 2005;65:6063–9. [PubMed: 16024606]
23. Ward RL, Cheong K, Ku SL, et al. Adverse prognostic effect of methylation in colorectal cancer is reversed by microsatellite instability. *J Clin Oncol* 2003;21:3729–36. [PubMed: 14551292]

24. Van Rijnsoever M, Elsaleh H, Joseph D, et al. CpG island methylator phenotype is an independent predictor of survival benefit from 5-fluorouracil in stage III colorectal cancer. *Clin Cancer Res* 2003;9:2898–903. [PubMed: 12912934]
25. Ogino S, Meyerhardt JA, Kawasaki T, et al. CpG island methylation, response to combination chemotherapy, and patient survival in advanced microsatellite stable colorectal carcinoma. *Virchows Arch* 2007;450:529–37. [PubMed: 17372756]
26. Shen L, Catalano PJ, Benson AB 3rd, et al. Association between DNA Methylation and Shortened Survival in Patients with Advanced Colorectal Cancer Treated with 5-Fluorouracil Based Chemotherapy. *Clin Cancer Res* 2007;13:6093–8. [PubMed: 17947473]
27. Lee S, Cho NY, Choi M, et al. Clinicopathological features of CpG island methylator phenotype-positive colorectal cancer and its adverse prognosis in relation to KRAS/BRAF mutation. *Pathol Int* 2008;58:104–13. [PubMed: 18199160]
28. Ferracin M, Gafa R, Miotto E, et al. The methylator phenotype in microsatellite stable colorectal cancers is characterized by a distinct gene expression profile. *J Pathol* 2008;214:594–602. [PubMed: 18241079]
29. Ogino S, Kawasaki T, Kirkner GJ, et al. Evaluation of markers for CpG island methylator phenotype (CIMP) in colorectal cancer by a large population-based sample. *J Mol Diagn* 2007;9:305–14. [PubMed: 17591929]
30. Colditz GA, Hankinson SE. The Nurses' Health Study: lifestyle and health among women. *Nat Rev Cancer* 2005;5:388–96. [PubMed: 15864280]
31. Chan AT, Ogino S, Fuchs CS. Aspirin and the Risk of Colorectal Cancer in Relation to the Expression of COX-2. *New Engl J Med* 2007;356:2131–42. [PubMed: 17522398]
32. Ogino S, Kawasaki T, Brahmandam M, et al. Sensitive sequencing method for KRAS mutation detection by Pyrosequencing. *J Mol Diagn* 2005;7:413–21. [PubMed: 16049314]
33. Ogino S, Kawasaki T, Kirkner GJ, et al. CpG island methylator phenotype-low (CIMP-low) in colorectal cancer: possible associations with male sex and KRAS mutations. *J Mol Diagn* 2006;8:582–8. [PubMed: 17065427]
34. Eads CA, Danenberg KD, Kawakami K, et al. MethyLight: a high-throughput assay to measure DNA methylation. *Nucleic Acids Res* 2000;28:E32. [PubMed: 10734209]
35. Ogino S, kawasaki T, Brahmandam M, et al. Precision and performance characteristics of bisulfite conversion and real-time PCR (MethyLight) for quantitative DNA methylation analysis. *J Mol Diagn* 2006;8:209–17. [PubMed: 16645207]
36. Kawasaki T, Ohnishi M, Nosho K, et al. CpG island methylator phenotype-low (CIMP-low) colorectal cancer shows not only fewer methylated CIMP-high-specific CpG island, but also low-level methylation in individual loci. *Mod Pathol* 2008;21:245–55. [PubMed: 18204436]
37. Boland CR, Thibodeau SN, Hamilton SR, et al. A National Cancer Institute Workshop on Microsatellite Instability for cancer detection and familial predisposition: development of international criteria for the determination of microsatellite instability in colorectal cancer. *Cancer Res* 1998;58:5248–57. [PubMed: 9823339]
38. Jass JR. Classification of colorectal cancer based on correlation of clinical, morphological and molecular features. *Histopathology* 2007;50:113–30. [PubMed: 17204026]
39. Ogino S, Goel A. Molecular classification and correlates in colorectal cancer. *J Mol Diagn* 2008;10:13–27. [PubMed: 18165277]
40. Deng G, Bell I, Crawley S, et al. BRAF mutation is frequently present in sporadic colorectal cancer with methylated hMLH1, but not in hereditary nonpolyposis colorectal cancer. *Clin Cancer Res* 2004;10:191–5. [PubMed: 14734469]
41. Domingo E, Laiho P, Ollikainen M, et al. BRAF screening as a low-cost effective strategy for simplifying HNPCC genetic testing. *J Med Genet* 2004;41:664–8. [PubMed: 15342696]
42. Young J, Jenkins M, Parry S, et al. Serrated pathway colorectal cancer in the population: genetic consideration. *Gut* 2007;56:1453–9. [PubMed: 17566021]
43. Le Gouvello S, Bastuji-Garin S, Aloulou N, et al. High prevalence of Foxp3 and IL-17 in MMR-proficient colorectal carcinomas. *Gut* 2008;57:772–9. [PubMed: 17965063]

44. Lees NP, Harrison KL, Hall CN, et al. Human colorectal mucosal O6-alkylguanine DNA-alkyltransferase activity and DNA-N7-methylguanine levels in colorectal adenoma cases and matched referents. *Gut* 2007;56:380–4. [PubMed: 16891355]
45. Ogino S, kawasaki T, Kirkner GJ, et al. Molecular correlates with MGMT promoter methylation and silencing support CpG island methylator phenotype-low (CIMP-low) in colorectal cancer. *Gut* 2007;56:1409–16. [PubMed: 17872570]
46. Boland CR. Clinical uses of microsatellite instability testing in colorectal cancer: an ongoing challenge. *J Clin Oncol* 2007;25:754–6. [PubMed: 17228016]
47. Morris EJ, Maughan NJ, Forman D, et al. Who to treat with adjuvant therapy in Dukes B/stage II colorectal cancer? The need for high quality pathology. *Gut* 2007;56:1419–25. [PubMed: 17494107]
48. Walther A, Houlston R, Tomlinson I. Association between chromosomal instability and prognosis in colorectal cancer: a meta-analysis. *Gut* 2008;57:941–50. [PubMed: 18364437]
49. Meyerhardt JA, Giovannucci EL, Holmes MD, et al. Physical activity and survival after colorectal cancer diagnosis. *J Clin Oncol* 2006;24:3527–34. [PubMed: 16822844]




Figure 1. Distribution of colon cancers and *BRAF* and *KRAS* mutations according to the number of methylated CIMP markers

A bimodal distribution of tumors is evident, and *BRAF* mutation is common in heavily methylated tumors, while *KRAS* mutation is common in tumors with fewer methylated markers. B(-), *BRAF* wild-type; CIMP, CpG island methylator phenotype; K(-), *KRAS* wild-type.




Figure 2. Kaplan-Meier survival curves in colon cancer according to combined CIMP/BRAF status
 A. Colon cancer-specific survival. B. Overall survival.
 CIMP-low/0 BRAF(+) tumors demonstrate shorter survival than the other 3 CIMP/BRAF types.
 P values indicate statistical significance of a deviation of any one of the curves from the null hypothesis.

Table 1
Clinical and molecular characteristics of colon cancer according to MSI or CIMP status.

| Clinical or molecular feature | All cases | MSI-high | MSS | CIMP-high(6-8 methylated promoters) | CIMP-low(1-5 methylated promoters) | CIMP-0(0 methylated promoters) |
|-------------------------------|----------------|----------------|----------------|-------------------------------------|------------------------------------|--------------------------------|
| Total N | 649 | 121 | 528 | 126 | 252 | 271 |
| Sex | | | | | | |
| Male (HPFS) | 283 (44%) | 43 (36%) | 240 (45%) | 40 (32%) | 125 (50%) | 118 (44%) |
| Female (NHS) | 366 (56%) | 78 (64%) | 288 (55%) | 86 (68%) | 127 (50%) | 153 (56%) |
| Mean age \pm SD | 66.5 \pm 8.2 | 68.1 \pm 7.3 | 66.2 \pm 8.4 | P=0.004 | 66.4 \pm 8.2 | 65.3 \pm 8.5 |
| Year of diagnosis | | | | P<0.0001 [#] * | | |
| Prior to 1990 | 101 (16%) | 13 (11%) | 88 (17%) | 9 (7.1%) | 43 (17%) | 49 (18%) |
| 1990 to 1999 | 473 (73%) | 95 (79%) | 378 (71%) | 102 (81%) | 184 (73%) | 187 (69%) |
| 2000 to 2002 | 75 (12%) | 13 (11%) | 62 (12%) | 15 (12%) | 25 (9.9%) | 35 (13%) |
| Tumor location [^] | | | | P=0.14 | | |
| Proximal | 371 (58%) | 102 (85%) | 269 (52%) | 114 (90%) | 153 (62%) | 104 (39%) |
| Distal | 269 (42%) | 18 (15%) | 251 (48%) | 12 (9.5%) | 94 (38%) | 163 (61%) |
| Tumor stage | | | | P<0.0001 | | |
| I | 137 (21%) | 19 (16%) | 118 (22%) | 14 (11%) | 55 (22%) | 68 (25%) |
| II | 216 (33%) | 70 (58%) | 146 (28%) | 66 (52%) | 64 (25%) | 86 (32%) |
| III | 165 (25%) | 22 (18%) | 143 (27%) | 24 (19%) | 69 (27%) | 72 (27%) |
| IV | 83 (13%) | 5 (4.1%) | 78 (15%) | 16 (13%) | 40 (16%) | 27 (10%) |
| Unknown | 48 (7.4%) | 5 (4.1%) | 43 (8.1%) | 6 (4.8%) | 24 (9.5%) | 18 (6.6%) |
| Tumor grade | | | | P<0.0001 | | |
| Low | 574 (89%) | 86 (71%) | 488 (93%) | 86 (68%) | 232 (92%) | 256 (95%) |
| High | 72 (11%) | 35 (29%) | 37 (7.0%) | 40 (32%) | 19 (7.6%) | 13 (4.8%) |
| KRAS mutation | | | | P<0.0001 | | |
| (-) | 409 (63%) | 102 (84%) | 307 (58%) | 109 (87%) | 131 (52%) | 169 (63%) |

| Clinical or molecular feature | All cases | MSI-high | MSS | CIMP-high(6-8 methylated promoters) | CIMP-low(1-5 methylated promoters) | CIMP-0(0 methylated promoters) |
|-------------------------------|-----------|----------------------|-----------|-------------------------------------|------------------------------------|--------------------------------|
| (+) | 238 (37%) | 19 (16%) P<0.0001 | 219 (42%) | 17 (13%) P<0.0001 | 120 (48%) | 101 (37%) |
| <i>BRAF</i> mutation | | | | | | |
| (-) | 526 (83%) | 65 (55%) | 461 (90%) | 49 (40%) | 221 (90%) | 256 (97%) |
| (+) | 105 (17%) | 53 (45%) P<0.0001 | 52 (10%) | 74 (60%) P<0.0001 | 24 (9.8%) | 7 (2.7%) |
| CIMP | | | | | | |
| CIMP-0 | 271 (42%) | 12 (9.9%) | 259 (49%) | - | - | - |
| CIMP-low | 252 (39%) | 21 (17%) | 231 (44%) | - | - | - |
| CIMP-high | 126 (19%) | 88 (73%) P<0.0001 | 38 (7.2%) | - | - | - |

%, indicates the proportion of tumors with a specific clinical or molecular feature in a given subtype.

[^] Proximal colon includes cecum to transverse colon, and distal colon includes splenic flexure to sigmoid colon.

[#]The t-test assuming unequal variances was used to compare mean ages.

^{*} Compared to combined CIMP-low/CIMP-0.

CIMP, CpG island methylator phenotype; HPFS, Health Professionals Follow-up Study; MSI, microsatellite instability; MSS, microsatellite stable; NHS, Nurses' Health Study; SD, standard deviation.

Table 2
Survival analysis on colon cancer patients according to molecular features in tumor

| Molecular feature | Total N | Colon cancer-specific mortality | | Overall mortality | | | |
|-------------------|-----------|---------------------------------|------------------------|---------------------|------------------------|--------------------------|------------------|
| | | Deaths/person-years | Univariate HR (95% CI) | Deaths/person-years | Univariate HR (95% CI) | Multivariate HR (95% CI) | |
| MSI | MSS | 528 | 149/4653 | 1 (referent) | 238/4653 | 1 (referent) | 1 (referent) |
| | MSI-high | 121 | 14/1101 | 0.38 (0.22–0.66) | 43/1101 | 0.75 (0.54–1.04) | 0.89 (0.58–1.38) |
| CIMP | CIMP-0 | 271 | 70/2556 | 1 (referent) | 106/2556 | 1 (referent) | 1 (referent) |
| | CIMP-low | 252 | 66/2183 | 1.09 (0.78–1.53) | 116/2183 | 1.27 (0.98–1.66) | 1.01 (0.77–1.33) |
| | CIMP-high | 126 | 27/1016 | 0.88 (0.57–1.38) | 59/1016 | 1.35 (0.98–1.85) | 0.78 (0.47–1.29) |
| KRAS mutation | (–) | 409 | 98/3604 | 1 (referent) | 170/3604 | 1 (referent) | 1 (referent) |
| | (+) | 238 | 65/2137 | 1.10 (0.81–1.51) | 111/2137 | 1.10 (0.86–1.40) | 0.98 (0.74–1.28) |
| BRAF mutation | (–) | 526 | 127/4821 | 1 (referent) | 229/4821 | 1 (referent) | 1 (referent) |
| | (+) | 105 | 36/775 | 1.73 (1.20–2.51) | 52/775 | 1.46 (1.08–1.98) | 1.20 (0.79–1.80) |

The multivariate analysis model includes age, year of diagnosis, sex, tumor location, stage, tumor grade, and statuses of MSI, CIMP, KRAS and BRAF.

CI, confidence interval; CIMP, CpG island methylator phenotype; HR, hazard ratio; MSI, microsatellite instability; MSS, microsatellite stable.

Table 3

Stage-specific hazard ratio (HR) for colon cancer-specific mortality in CIMP-high tumors compared to CIMP-0 tumors.

| AJCC tumor stage | Total N | Multivariate HR (with 95% CI) for colon cancer- specific mortality in CIMP-high tumors compared to CIMP-0 tumors |
|------------------|---------|--|
| Stage I | 82 | 0* |
| Stage II | 152 | 0.76 (0.20–2.82) |
| Stage III | 96 | 0.52 (0.17–1.59) |
| Stage IV | 43 | 0.47 (0.18–1.21) |
| Stage missing | 24 | 0* |

The multivariate analysis model includes the CIMP variable (CIMP-high vs. CIMP-0) stratified by stage, age, year of diagnosis, sex, tumor location, tumor stage, grade, MSI, CIMP, *KRAS* and *BRAF*.

* 95% CI was not shown because there was no death in patients with stage I and stage-missing CIMP-high tumors.

AJCC, American Joint Commission on Cancer; CI, confidence interval; CIMP, CpG island methylator phenotype; HR, hazard ratio.

Table 4
 Combined MSI/CIMP status, MSI/*BRAF* status or CIMP/*BRAF* status and patient survival in colon cancer

| Molecular feature | N | Colon cancer-specific mortality | | Overall mortality | | Multivariate HR (95% CI) |
|----------------------------|-----|---------------------------------|--------------------------|---------------------|--------------------------|--------------------------|
| | | Deaths/person-years | Multivariate HR (95% CI) | Deaths/person-years | Multivariate HR (95% CI) | |
| MSS CIMP-low/0 | 490 | 134/4384 | 1 (referent) | 218/4384 | 1 (referent) | 1 (referent) |
| MSS CIMP-high | 38 | 15/269 | 0.40 (0.20–0.83) | 20/269 | 0.62 (0.34–1.13) | 0.62 (0.34–1.13) |
| MSI-high CIMP-low/0 | 33 | 2/355 | 0.17 (0.02–1.22) | 4/355 | 0.22 (0.07–0.69) | 0.22 (0.07–0.69) |
| MSI-high CIMP-high | 88 | 12/746 | 0.30 (0.14–0.64) | 39/746 | 0.77 (0.48–1.24) | 0.77 (0.48–1.24) |
| MSS <i>BRAF</i> (+) | 52 | 25/365 | 1 (referent) | 28/365 | 1 (referent) | 1 (referent) |
| MSS <i>BRAF</i> (-) | 461 | 121/4162 | 0.74 (0.41–1.36) | 200/4162 | 1.12 (0.67–1.89) | 1.12 (0.67–1.89) |
| MSI-high <i>BRAF</i> (+) | 53 | 11/410 | 1.09 (0.48–2.51) | 24/410 | 1.27 (0.68–2.36) | 1.27 (0.68–2.36) |
| MSI-high <i>BRAF</i> (-) | 65 | 3/659 | 0.18 (0.05–0.61) | 18/659 | 0.65 (0.35–1.21) | 0.65 (0.35–1.21) |
| CIMP-low/0 <i>BRAF</i> (+) | 31 | 17/199 | 1 (referent) | 20/199 | 1 (referent) | 1 (referent) |
| CIMP-low/0 <i>BRAF</i> (-) | 477 | 116/4410 | 0.46 (0.25–0.86) | 193/4410 | 0.56 (0.33–0.95) | 0.56 (0.33–0.95) |
| CIMP-high <i>BRAF</i> (+) | 74 | 19/577 | 0.35 (0.16–0.79) | 32/577 | 0.49 (0.25–0.98) | 0.49 (0.25–0.98) |
| CIMP-high <i>BRAF</i> (-) | 49 | 8/411 | 0.30 (0.11–0.82) | 25/411 | 0.63 (0.30–1.31) | 0.63 (0.30–1.31) |

The multivariate analysis model includes age, year of diagnosis, sex, tumor location, stage, tumor grade, *KRAS* status, as well as *BRAF* status in the top analysis (for MSI/CIMP types), CIMP status in the middle analysis (for MSI/*BRAF* types) or MSI status in the bottom analysis (for CIMP/*BRAF* types).

CI, confidence interval; CIMP, CpG island methylator phenotype; HR, hazard ratio; MSI, microsatellite instability; MSS, microsatellite stable.